

Vortices on Higher Genus Surfaces[★]Kai-Ming Lee[†]*California Institute of Technology, Pasadena, CA 91125***Abstract**

We consider the topological interactions of vortices on general surfaces. If the genus of the surface is greater than zero, the handles can carry magnetic flux. The classical state of the vortices and the handles can be described by a mapping from the fundamental group to the unbroken gauge group. The allowed configurations must satisfy a relation induced by the fundamental group. Upon quantization, the handles can carry “Cheshire charge.” The motion of the vortices can be described by the braid group of the surface. How the motion of the vortices affects the state is analyzed in detail.

★ Work supported in part by the U.S. Dept. of Energy under Grant no. DE-FG03-92-ER40701.

† email address: kmlee@theory3.caltech.edu

1. Introduction

When a gauge symmetry is spontaneously broken, in general there will be stable topological defects.^[1] What types of defects will be created depends on the spacetime dimension and the topology of the vacuum manifold. In two spatial dimensions, if the fundamental group of the vacuum manifold is non-trivial, there will be point defects which are called vortices. A charged particle winding around a vortex will be transformed by an element of the unbroken gauge group. This is the (non-abelian) Aharonov-Bohm effect.^[2–12] It is long-range and topological. This means that the gauge transformation will not depend either on how far apart the particle and the vortex are, or on the exact loops the charged particle travels along, as long as their linking numbers with the vortex are the same. We will say that the vortex carries (non-abelian) magnetic flux.

Another way to look at it is that in the presence of the vortices, the fundamental group of the surface is non-trivial.^[7] After a charged particle travels along a non-contractible loop around a vortex, it will remain the same only up to a gauge transformation. The element of the unbroken gauge group associated with that transformation is the magnetic flux carried by the vortex. However, the fundamental group of the surface may be non-trivial even without any vortices. For example, there may be handles on the surface. There are two non-equivalent non-contractible loops associated to each handle. Then, by the same argument, we expect we can assign group elements to the two loops and the handle can carry magnetic flux; therefore, the handles will have topological interactions with the vortices and the charged particles. If we interchange two vortices or let a vortex go along a non-contractible loop, the magnetic flux carried by the vortices and the loop will be changed. This kind of motion of the vortices can be described by the braid group of the surface.^[13,14] We then have a natural action of the elements of the braid group on the states of the vortices and the surface.

If the surface is compact, there is one relation between the generators of the fundamental group of the surface, the group elements assigned to the vortices and

the non-contractible loops must satisfy a relation induced from that relation. This restricts the possible magnetic flux carried by the vortices and handles on any compact surfaces. The simplest example is that we cannot put a single non-trivial vortex on a sphere.

In the semiclassical approximation, a pair consisting of a vortex and an anti-vortex may carry electric charge, “Cheshire charge.”^[5,6,10] It turns out that the properties of a handle are similar (but not equal) to the properties of two vortex-anti-vortex pairs. In particular, a handle can carry Cheshire charge. If the size of the handle is very small, an observer outside the handle will see a “particle” that carries both magnetic flux and electric charge, a dyon.^[1] (The term “dyon” is originally for a particle that carries both electric and magnetic charge in $3+1$ dimensions. We stretch its meaning to $2+1$ dimensional spacetime.) In fact, any particle that carries magnetic flux and/or electric charge falls into the representations of the quantum double associated with the gauge group.^[15,16] In the language of the quantum double, we have a unified treatment of the magnetic flux and electric charge. There is also a restriction on the configurations of the dyons on any compact surface.

In section 2, the basic properties of vortices will be briefly reviewed. The purpose of this section is to establish conventions. In section 3, the braid group of a surface will be described and the topological interactions between vortices and handles will be analyzed. We will find out that, locally, there is no restriction on the assignment of group elements to the non-trivial loops of a handle. In section 4, we will give a semiclassical analysis of the theory. The argument that the handle can carry Cheshire charge is given. We also explain what a quantum double is and why it is relevant. In section 5, the most general formulation of dyons on a surface is described. (The analysis of the previous sections is a special case in this formulation.) We give the conclusions and some comments in section 6.

2. Non-abelian vortices

We assume that for our theory, in the high energy regime, the gauge group is a simply connected Lie group. In the low energy regime, the symmetry is spontaneously broken by the Higgs mechanism, say, to a finite group, G . Then in $2 + 1$ dimensions, the point defects will be classified by $\pi_0(G) \cong G$.^[1] From now on, we only consider the unbroken gauge group and its representations. The original broken gauge group plays no role in the following discussion. If the energy scale of the symmetry breaking is very high, the size of the vortices will be very small. Low energy experiment usually cannot probe the core of vortices. Then the space that a low energy particle sees is the original space with the points where the vortices are removed.

We can assign a group element to any isolated vortex to label the flux by the following method. Choose a fixed but arbitrary base point, x_0 (away from the vortex), and a loop around it. Then, calculate the untraced Wilson loop,

$$a(C, x_0) = P \exp \left(i \int_{C, x_0} A \cdot dx \right) , \quad (2.1)$$

where P denotes the path ordering. The orientation of the loop, C , is only a convention. We adopt the convention indicated in Fig. 1. Then, if a charged particle in representation (ν) of G is transported along the loop C , it will be transformed by $D^{(\nu)}(a(C, x_0))$:^[6]

$$v^{(\nu)} \rightarrow D^{(\nu)}(a(C, x_0))v^{(\nu)} , \quad (2.2)$$

where $v^{(\nu)}$ denotes the state of the charged particle.

Since the unbroken gauge group is discrete, there is no local low energy gauge excitation. The group element $a(C, x_0)$ is invariant under a continuous deformation of the loop C . This is how the fundamental group of the space comes in. We have assigned a group element to a generator of the fundamental group of the space with punctures.




Fig. 1: The vortex in this figure is represented by a circle with a cross. We choose a base point, x_0 , and a standard path, C , around the vortex to measure its flux.

If there are two or more vortices, we have to choose a standard loop for each vortex as in Fig. 2.^[7] Then we can assign group elements to the loops. The combined magnetic flux, for example, of vortex 1 and vortex 2 is the product of the group elements associated to them. Here we adopt the convention that the product of two loops, C_1C_2 , in the fundamental group means that the particle will travel C_2 first and then C_1 . So, the combined magnetic flux, in this convention, is $a(C_1C_2, x_0) = a(C_1, x_0)a(C_2, x_0)$.




Fig. 2: For two or more vortices, we choose one standard path for each vortex.

Let us consider what will happen if we interchange two vortices. Let the flux of vortex 1 and 2 be h_1 and h_2 respectively. If we interchange the vortices counter-clockwise, Fig. 3a, the magnetic flux of them will change. We have to find two loops

such that *after* the interchange, they will deform to the standard loops. Then the group elements associated with them are the magnetic flux of the vortices after the interchange. From Fig. 3b, we find that

$$\begin{aligned} h_1 &\rightarrow h_1 h_2 h_1^{-1} \\ h_2 &\rightarrow h_1 . \end{aligned} \tag{2.3}$$

We will rely on this kind of loop tracing method to calculate various processes in the coming sections.




Fig. 3: In a), the two paths are the standard paths based on the same based point. The dark curves with arrows represent the interchange of the two vortices. In b), the two paths will deform to the standard paths after the interchange of the two vortices. So the flux associated with them are the flux of the two vortices after the interchange.

3. Vortices on higher genus surfaces

The basic element of an orientable surface with genus greater than zero is a handle.^[17] All compact surfaces can be classified according to the number of handles they have. For a single handle, there are two generators in the fundamental group. We can choose the generators to be the loops α and β in Fig. 4. Then it is easy to see that the loop in Fig. 5 is equal to $\alpha\beta\alpha^{-1}\beta^{-1}$.




Fig. 4: The wide curves represent a handle that stands out of the paper. The two standard paths of the handle are the α in a) and β in b). Part of β is under the handle.




Fig. 5: We can use this path to calculate the flux of a handle. The path is equal to $\alpha\beta\alpha^{-1}\beta^{-1}$.

Since α and β are non-contractible, a charged particle transported along them may be transformed by an element of the gauge group. By sending charged particles along the loops, we can measure the group elements associated with them, for example,

$$\begin{aligned}\alpha &\mapsto a \\ \beta &\mapsto b\end{aligned}\tag{3.1}$$

where $a, b \in G$.

What will happen if a vortex winds around the loops? We expect that the magnetic flux of the vortex and group elements associated with the loops will change. We can calculate the changes by the loop tracing method as in section 2. This means that we have to find loops such that after the traveling of the vortex, these loops deform to the standard loops we used to measure the flux.

If the vortex with flux h winds around the loop in Fig. 6a, we will say that it winds around α . It implicitly means that we have chosen a path (in this case, the path can be a straight line segment,) from the position of the vortex to the base point and the vortex goes along this path, then the α defined in Fig. 4 and finally that path again in reverse. If also the elements associated with α and β are a and b respectively, from Fig. 6b, c and d, we find that they will change to

$$\begin{aligned}h &\rightarrow aha^{-1} \\ a &\rightarrow ahah^{-1}a^{-1} \\ b &\rightarrow aha^{-1}h^{-1}bah^{-1}a^{-1} .\end{aligned}\tag{3.2}$$

If the vortex winds around β in a similar sense, from Fig. 7, they will change to

$$\begin{aligned}h &\rightarrow h^{-1}bhb^{-1}h \\ a &\rightarrow ah \\ b &\rightarrow h^{-1}bh .\end{aligned}\tag{3.3}$$




Fig. 6: The motion of the vortex is represented by the loop in a). It is also called the α loop. After the motion of the vortex as in a), the path in b) will deform to the standard path in Fig. 1. The path in c) will deform to the standard path α in Fig. 4a. The path in d) will deform to the standard path β in Fig. 4b.

Is it possible to assign arbitrary group elements to α and β ? The answer is yes, at least locally. From (3.2) and (3.3), it is easy to see that even if the group is abelian and initially the element associated with each loop is the identity, after the winding of a non-trivial vortex, the group elements are no longer trivial. So, we can transfer the magnetic flux from a vortex to the handle by sending the vortex to go along α or β . To excite the handle to a state with $\alpha \mapsto a$ and $\beta \mapsto b$, consider the following: We assume that vortices with arbitrary flux exist and initially the group elements




Fig. 7: Similar to Fig. 6, the loop in a) represents the motion of the vortex. Paths in b), c) and d) will deform to the standard paths in Fig. 1, α and β in Fig. 4 respectively.

associated with α and β are the identity. And we send an $ab^{-1}a^{-1}$ vortex to go along α . Then, we send an a vortex to go along β . By (3.2) and (3.3), we have

$$\begin{array}{ccccccc} \alpha & \mapsto & e & \xrightarrow{\alpha} & e & \xrightarrow{\beta} & a \\ \beta & \mapsto & e & \longrightarrow & aba^{-1} & \longrightarrow & b \end{array} . \quad (3.4)$$

Thus, after the vortices execute the prescribed motion, the loops α and β are associated with the desired group elements a and b respectively.

If the throat of the handle is small or we ignore the internal structure of the

handle and there are no pointlike vortices hiding inside the handle, we can measure the flux of the “particle” by the loop in Fig. 5. The flux is $aba^{-1}b^{-1}$; the flux carried by a handle must be in this form.

Now let us formulate the theory in precise mathematical terms. Let the space be an orientable connected surface, Σ . If there are n vortices on it, we have to consider the fundamental group of the surface with n punctures and a base point, $\pi_1(\Sigma(n), x_0)$. The combined magnetic flux (or group elements) of vortices or handles follows from the multiplication rule of the fundamental group. Any classical state of the vortices and the surface is a homomorphism from $\pi_1(\Sigma(n), x_0)$ to G ,

$$\rho : \pi_1(\Sigma(n), x_0) \rightarrow G . \quad (3.5)$$

If the surface is also compact, there is one relation between the generators of the fundamental group. For example, the relation for a surface of genus g , Σ_g , is^[17]

$$\alpha_1\beta_1\alpha_1^{-1}\beta_1^{-1} \dots \alpha_g\beta_g\alpha_g^{-1}\beta_g^{-1} = e . \quad (3.6)$$

For the compact surface of genus g with n punctures, $\Sigma_g(n)$, the relation is

$$\alpha_1\beta_1\alpha_1^{-1}\beta_1^{-1} \dots \alpha_g\beta_g\alpha_g^{-1}\beta_g^{-1}C_1 \dots C_n = e \quad (3.7)$$

where our convention for the loops is shown in Fig. 8.

The flux of the vortices and handles must satisfy this relation. For example, the relation associated with a single vortex on a sphere is

$$C_1 = e . \quad (3.8)$$

It is inconsistent to put a single non-trivial vortex on a sphere. Also, the relation of a torus without any vortex is

$$\alpha_1\beta_1\alpha_1^{-1}\beta_1^{-1} = e . \quad (3.9)$$

Therefore, the group elements associated with the two loops must commute.




Fig. 8: A compact surface is represented by a large sphere with handles here. We choose the standard positions of the n vortices and the g handles as in this figure. We also choose a standard path for each vortex and the two standard paths α and β for each handle. The positions and the paths are consistent with Fig. 2 and Fig. 4.

This is a global constraint on the possible flux. We have seen that we can excite the handle to any state locally. We will show that it is possible to construct the state corresponding to any homomorphism, $\rho : \pi_1(\Sigma(n), x_0) \rightarrow G$. We assume that we can create vortex-anti-vortex pairs with arbitrary flux; to construct the state corresponding to ρ , we create two sets of vortex-anti-vortex pairs. The first set consists of n pairs. They contain exactly the n vortices that we want. Then we have n anti-vortices left and push the anti-vortices to some simply-connected region. The second set of vortex-anti-vortex pairs contains $2g$ pairs with appropriate flux. By sending them to go along the α 's and the β 's, we can excite the handles to the desired states. After they go along the loops, their flux will be changed. Now, the combined magnetic flux of the resulting second set of $4g$ vortices will no longer be trivial. Let us push them to the same simply-connected region that contains the n anti-vortices. We claim that the combined magnetic flux of the first set of anti-vortices and the second set of vortices is trivial. Since the surface is compact, a loop wrapped around that region can be deformed to a loop that wraps around all handles and the n vortices.

The magnetic flux measured along this loop must be trivial because the flux assigned by ρ satisfies the relation (3.7). This means that the combined magnetic flux of the left-over vortices is trivial. They can completely annihilate each other and the state of the surface with the n vortices is given by ρ . We conclude that the space of all states is $\text{Hom}(\pi_1(\Sigma(n), x_0), G)$.

When we consider the kinematics of the vortices on the surface, at low energy, they are not allowed to collide with each other. And because the magnetic flux we measure depends on the loops we choose, to determine the flux after any motion, the vortices must be brought back to some standard positions. This kind of motion is exactly described by the braid group of the surface.

If a collision of particles is not allowed, the configuration space of n distinguishable particles on a surface, Σ , is $\Sigma^n - \Delta$ where Δ is the subset of Σ^n in which at least two points in the Cartesian product coincide. The permutation group S_n has an obvious action on this configuration space. The configuration space of n indistinguishable particles is then $(\Sigma^n - \Delta)/S_n$. The definition of the braid group of n points on the surface is^[13,14]

$$B_n(\Sigma) = \pi_1((\Sigma^n - \Delta)/S_n) . \quad (3.10)$$

If the surface is the plane, R^2 , the braid group $B_n(R^2)$ is the usual braid group with $n - 1$ generators which interchange adjacent points. In $B_n(\Sigma_g)$, there are $2g$ more generators. They are the α_i and β_i , $1 \leq i \leq g$ as shown in Fig. 9.[★] (We use the same symbols α and β to denote the loops in the fundamental group and the generators of the braid group as explained above (3.2).)

These generators are not independent. They satisfy, in our convention, the fol-

★ We use a different convention from Ref. 13 because the α 's there involve all handles.

lowing relations.

$$\begin{aligned}
\sigma_i \sigma_j &= \sigma_j \sigma_i & |i - j| \geq 2, \\
\sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1} & 1 \leq i \leq n - 2, \\
\alpha_1 \beta_1 \alpha_1^{-1} \beta_1^{-1} \dots \alpha_g \beta_g \alpha_g^{-1} \beta_g^{-1} \sigma_{n-1} \dots \sigma_1^2 \dots \sigma_{n-1} &= e, \\
\sigma_i \alpha_l \sigma_i^{-1} \alpha_l^{-1} &= e & 2 \leq i \leq n - 1, \quad 1 \leq l \leq g, \\
\sigma_i \beta_l \sigma_i^{-1} \beta_l^{-1} &= e & 2 \leq i \leq n - 1, \quad 1 \leq l \leq g, \\
\sigma_1 \alpha_p \sigma_1^{-1} \alpha_l &= \alpha_l \sigma_1 \alpha_p \sigma_1^{-1} & 1 \leq p < l \leq g, \\
\sigma_1 \beta_p \sigma_1^{-1} \beta_l &= \beta_l \sigma_1 \beta_p \sigma_1^{-1} & 1 \leq p < l \leq g, \\
\sigma_1 \alpha_p \sigma_1 \alpha_p &= \alpha_p \sigma_1 \alpha_p \sigma_1 & 1 \leq p \leq g, \\
\sigma_1 \beta_p \sigma_1 \beta_p &= \beta_p \sigma_1 \beta_p \sigma_1 & 1 \leq p \leq g, \\
\alpha_p \sigma_1^{-1} \beta_l \sigma_1 &= \sigma_1^{-1} \beta_l \sigma_1 \alpha_p & 1 \leq p < l \leq g, \\
\beta_p \sigma_1^{-1} \alpha_l \sigma_1 &= \sigma_1^{-1} \alpha_l \sigma_1 \beta_p & 1 \leq p < l \leq g, \\
\sigma_1 \alpha_p \sigma_1 \beta_p &= \beta_p \sigma_1 \alpha_p \sigma_1^{-1} & 1 \leq p \leq g.
\end{aligned} \tag{3.11}$$




Fig. 9: These are the conventions for the braidings of the vortices with each other and with the handles. They are equal to the braidings in Fig. 3a, Fig. 6a and Fig. 7a.

There is a natural action of the braid group of surface with n points on the fundamental group of the surface with n punctures defined as follow. If $\tau \in B_n(\Sigma)$ and $\gamma \in \pi_1(\Sigma(n), x_0)$, define $\tau(\gamma)$ to be the loop such that *after* the motion of the n points according to τ , the loop will deform to γ . By the calculation in (2.3), (3.2) and (3.3), we find that the nontrivial actions for $B_n(\Sigma_g)$ are

$$\begin{aligned}
\sigma_i(C_i) &= C_{i+1} \\
\sigma_i(C_{i+1}) &= C_{i+1}C_iC_{i+1}^{-1} \\
\alpha_j(C_1) &= \alpha_jC_1\alpha_j^{-1} \\
\alpha_j(\alpha_j) &= \alpha_jC_1\alpha_jC_1^{-1}\alpha_j^{-1} \\
\alpha_j(\beta_j) &= \alpha_jC_1\alpha_j^{-1}C_1^{-1}\beta_j\alpha_jC_1^{-1}\alpha_j^{-1} \\
\beta_j(C_1) &= C_1^{-1}\beta_jC_1\beta_j^{-1}C_1 \\
\beta_j(\alpha_j) &= \alpha_jC_1 \\
\beta_j(\beta_j) &= C_1^{-1}\beta_jC_1 .
\end{aligned} \tag{3.12}$$

It is easy to check that this definition satisfies the relations (3.11). This action induces an action of the braid group on the states of the vortices and surface. And this is exactly how the state will be changed after the motion of the vortices.

4. Semi-classical analysis

We are going to argue that if we specify the flux of α and β of a handle, we know the quantum state of the handle completely. (Of course, a general quantum state of the handle could be a linear combination of the flux eigenstates of α and β .) The scheme is as follows. We try to find out a complete set of commuting observables by first choosing an observable, say A , and find out its eigenstates. In general, there is more than one independent eigenvector with the same eigenvalue. So, we find another observable, B , which commutes with A . Then we can decompose the eigenspaces of A with respect to B . If the dimensions of the simultaneous eigenspaces of A and B are still greater than one, we find yet another observable which commutes with both

A and B and decompose the eigenspaces and so on. This process will stop if all the simultaneous eigenspaces are one-dimensional or we run out of observables.

In our case of discrete gauge theory, there are not many observables. First of all, the theory is topological. We don't have any local excitations, and the only things that we can measure are the magnetic flux and the electric charge bounded by a loop. For a handle, we can send charged particles to go along α and β to measure the flux of them. The measurement of one loop does not affect the flux of the other, therefore these two observables commute. Let us denote the state of a handle that α maps to a and β maps to b by $|a, b, X\rangle$ where X specifies any other quantum numbers needed to completely specify the state of a handle. Our objective is to prove that no such X is needed. Now, the only other possible degrees of freedom, X , are the charge bounded by the two loops. It turns out that we cannot measure the charge bounded by α , say, without messing up the flux of β . The charge measurement of α does not commute with the measurement of the flux of β and vice versa. Since these are all the observables in the theory, the flux of α and β form a complete set of commuting observables and we do not need any X .

We now explain why we cannot measure the charge bounded by α without affecting the flux of β . The only way we can measure the charge bounded by a loop is to send vortices along the loop and deduce the charge from the interference pattern.^[8] (We explain how to measure flux of a vortex by charged particles and how to measure the charge of a particle by vortices in the Appendix.) If the handle is in the state $|a, b, X\rangle$ and the flux of the vortex is h , the initial state is $|h\rangle \otimes |a, b, X\rangle$. Now, suppose that the vortex winds around the loop α , from (3.2), the final state is $|aha^{-1}\rangle \otimes |ahah^{-1}a^{-1}, aha^{-1}h^{-1}bah^{-1}a^{-1}, X'\rangle$ where the quantum number X may change to X' after the winding of the vortex. The interference term is $\langle h|aha^{-1}\rangle \langle a, b, X|ahah^{-1}a^{-1}, aha^{-1}h^{-1}bah^{-1}a^{-1}, X'\rangle$. The first factor is non-zero if a and h commute, but then the second factor is $\langle a, b, X|a, bh^{-1}, X'\rangle$ which is zero for non-trivial h ; there is no interference and we cannot know the charge bounded by α .

Notice that in some state of the handle, α can bound some well-defined electric charge. For example, in the state $\frac{1}{\sqrt{|G|}} \sum_{b \in G} |e, b\rangle$, the charge bounded by α can be measured and it is trivial.[★] Some other linear combinations will give non-trivial charge, however, none of these α -charge eigenstates are β -flux eigenstates.

Recall that if the state of the handle is $|a, b\rangle$, it carries magnetic flux $aba^{-1}b^{-1}$. In the neighborhood of such a handle, (in general, in the neighborhood of a vortex with non-trivial magnetic flux), it is impossible to implement a global gauge transformation h if h does not commute with the flux of the handle. That is when we try to extend a local gauge transformation h along a loop around the handle, the transformation will be conjugated by the flux of the handle at the end of the loop. There is no way to solve this inconsistent boundary condition, called the global color problem.^[18] We can only consider global gauge transformations that are in the normalizer of $aba^{-1}b^{-1}$, $N_{aba^{-1}b^{-1}}$. Under such a gauge transformation h , the state is transformed to

$$|a, b\rangle \rightarrow |hah^{-1}, hbh^{-1}\rangle . \quad (4.1)$$

Semi-classically, linear combinations of these states are physically attainable. The vector space spanned by all these states of a handle can be decomposed to a direct sum of irreducible representations of $N_{aba^{-1}b^{-1}}$. These irreducible representations are the possible Cheshire charges that a handle can carry. Notice that the mathematical structure of this vector space is equal to the structure of the states of two vortex-anti-vortex pairs. They can carry the same kinds of Cheshire charge.

Particles that carry both magnetic flux and electric charge are called dyons. The mathematical tool to classify them is the quantum double of a group and its representations. We will give a brief review of the necessary details here. Interested readers can look up the references for a full account.^[15,16,19]

The difficulty of classifying dyons is that when a dyon carries magnetic flux a , we can only consider electric charges which fall into the representations of the normalizer

★ I thank Hoi-Kwong Lo for pointing this out to me.

of a , N_a . If there are two dyons with flux a and b , their electric charges will be classified by N_a and N_b respectively. However, when we consider the two dyons as a whole, the total magnetic flux will be ab (in some convention) and the electric charge must be a representation of N_{ab} . We will find out that the irreducible representations of the quantum double have exactly this property. They are labeled by the flux and an irreducible representation of the normalizer of the flux. A tensor product of two irreducible representations can be decomposed to a direct sum of irreducible representations of the normalizer of the total flux. There is also an element in (the tensor product of two copies of) the quantum double to implement the braiding operation.

Let us begin by recalling some properties of representations of a group. Any representation of a group, G , on a vector space, V , is a homomorphism

$$\phi : G \rightarrow \text{End}(V) . \quad (4.2)$$

This homomorphism can be extended linearly to the group algebra, $\mathcal{C}[G]$, by

$$\phi(\sum k_i h_i) = \sum k_i \phi(h_i) , \quad (4.3)$$

where $k_i \in \mathcal{C}$. When we consider the tensor product of two representations, $\phi = \phi_1 \times \phi_2$, we have

$$\phi(h) = \phi_1(h) \otimes \phi_2(h) , \quad (4.4)$$

if h is a group element. In order to lift to the group algebra, we define the comultiplication, $\Delta : \mathcal{C}[G] \rightarrow \mathcal{C}[G] \otimes \mathcal{C}[G]$, by

$$\Delta(\sum k_i h_i) = \sum k_i h_i \otimes h_i . \quad (4.5)$$

Then, $\phi(h) = (\phi_1 \otimes \phi_2)\Delta(h)$ where now h can be any element in the group algebra.

The physical meaning of the comultiplication is that when a system consists of two subsystems, comultiplication bridges between the transformation of the whole system and the individual transformations of the subsystems. In general, if the symmetry transformations of a theory form an algebra, we expect there is a corresponding comultiplication to relate the symmetry transformations of the whole system and the subsystems.

Now, consider the gauge theory of a finite group, G , in two spatial dimensions. From the above discussion, we expect to have the following operators. For each element, a , of G , there is the gauge transformation operator of a . We can implement this operator by sending an a vortex around the base point (in some convention). The system does not change but the basis we used to measure the flux and charge has changed by a gauge transformation. It is equivalent to relabel everything in the system. For example, if the flux of a vortex is initially labeled by h , after the transformation, it is labeled by aha^{-1} . We denote this operator by the same symbol, a .

An observer far away from the system can also measure the total magnetic flux of the system relative to some fixed gauge choice. We also have a projection operator, P_a , for each $a \in G$, to project to the subspace of the total flux, a . The algebra of operators, $D(G)$, is generated by a and P_b where a and $b \in G$.

The multiplication of a , b in $D(G)$ is same as the multiplication in the group. Since P_a is a projection operator,

$$P_a P_b = \delta_{ab} P_a . \quad (4.6)$$

After a gauge transformation of a , the magnetic flux of the system changes from b to aba^{-1} , giving us

$$a P_b = P_{aba^{-1}} a . \quad (4.7)$$

We have completely determined the algebraic structure of $D(G)$.

The comultiplication, $\Delta : D(G) \rightarrow D(G) \otimes D(G)$, of elements of G is the one we discussed before

$$\Delta(a) = a \otimes a . \quad (4.8)$$

If the system is composed of two subsystems and the magnetic flux of them is b and c , then the flux of the whole system is bc . Conversely, if the total flux is a , the flux of the two subsystems can be any b and c as long as $bc = a$, giving us

$$\Delta(P_a) = \sum_{\substack{b,c \\ bc=a}} P_b \otimes P_c . \quad (4.9)$$

In this equation, we have implicitly assumed some standard paths are chosen. Then for general elements in $D(G)$, the comultiplication is[★]

$$\begin{aligned} \Delta(P_a b) &= \Delta(P_a) \Delta(b) \\ &= \left(\sum_{\substack{c,d \\ cd=a}} P_c \otimes P_d \right) (b \otimes b) \\ &= \sum_{\substack{c,d \\ cd=a}} (P_c b) \otimes (P_d b) . \end{aligned} \quad (4.10)$$




Fig. 10: This is essentially the same as Fig. 3. The shaded areas are the locations of the subsystems. Region X and Y are bounded by dotted lines.

★ $P_a b$ in Ref. 15 and 16 is written as $a \underset{b}{\perp}$.

Let us consider the two subsystems, S_1 and S_2 , located in region X and region Y respectively. Our convention is that the first factor in $D(G) \otimes D(G)$ acts on the system in region X and the second factor acts on system in region Y . What will happen if the two subsystems interchange positions as in Fig. 10? If the magnetic flux of S_1 is b , the effect of the braiding is that apart from the position change, S_1 does not change its state but S_2 will be changed by a gauge transformation b ,

$$|S_1\rangle \otimes |S_2\rangle \rightarrow (b|S_2\rangle) \otimes |S_1\rangle . \quad (4.11)$$

If S_1 is not in a magnetic flux eigenstate, we have

$$|S_1\rangle \otimes |S_2\rangle \rightarrow \sum_{b \in G} (b|S_2\rangle) \otimes (P_b|S_1\rangle) . \quad (4.12)$$

If we define τ to be the operator to interchange the two factors in a tensor product and $R = \sum_b P_b \otimes b \in D(G) \otimes D(G)$, then the above action can be described by an operator $\mathcal{R} = \tau \circ R$ because

$$\begin{aligned} \mathcal{R}(|S_1\rangle \otimes |S_2\rangle) &= \tau\left(\sum_{b \in G} (P_b|S_1\rangle) \otimes (b|S_2\rangle)\right) \\ &= \sum_{b \in G} (b|S_2\rangle) \otimes (P_b|S_1\rangle) . \end{aligned} \quad (4.13)$$

It is easy to show that $R^{-1} = \sum_b P_b \otimes b^{-1}$ and for any $P_a b \in D(G)$,

$$\begin{aligned}
R\Delta(P_ab)R^{-1} &= R(\Delta P_a)(\Delta b)R^{-1} \\
&= \sum_{\substack{c,d \\ hk=a}} (P_c \otimes c)(P_h \otimes P_k)(b \otimes b)(P_d \otimes d^{-1}) \\
&= \sum P_c P_h b P_d \otimes c P_k b d^{-1} \\
&= \sum P_c P_h P_{b d b^{-1}} b \otimes c P_k b d^{-1} \\
&= \sum_{hk=a} P_h b \otimes h P_k b (b^{-1} h^{-1} b) \\
&= \sum_{hk=a} P_h b \otimes P_{h k h^{-1}} b \\
&= \sum_{hk=a} P_k b \otimes P_h b \\
&= \tau(\Delta(P_a b)) .
\end{aligned} \tag{4.14}$$

The meaning of this equation is the following. If an operator $d \in D(G)$ acts on the whole system, we can calculate its effect on the subsystems either by directly applying the comultiplication or by the following procedure. First, interchange the two subsystems in clockwise direction. Then, apply the comultiplication and finally, interchange the subsystems (counterclockwise) back to their original positions.

With the multiplication, comultiplication and the R operator, (and some other structures) the algebra $D(G)$ is called the quantum double associated with the group G .^[20] We have seen that the quantum double is a generalization of the group algebra and it has direct physical meaning in a theory with vortices. The particles in such a theory will fall into representations of the quantum double.

Now we describe the irreducible representations of $D(G)$.^[15] Let the set of all conjugacy classes of G be $\{{}^A C\}$. The conjugacy class containing a will be denoted by $[a]$. For each class, fix an ordering of the elements $\{{}^A C\} = \{{}^A g_1, \dots, {}^A g_k\}$. Let ${}^A N$ be the normalizer of ${}^A g_1$. Choose elements, ${}^A x_1, \dots, {}^A x_k \in G$, such that ${}^A g_i = {}^A x_i {}^A g_1 {}^A x_i^{-1}$. We take ${}^A x_1 = e$. Consider the vector space, V_ν^A , spanned by the vectors $|{}^A g_j, {}^\nu v_i\rangle$, $j = 1, \dots, k$ and $i = 1, \dots, \dim \nu$, where $\{{}^\nu v_i\}$ is a basis of the ν irreducible representation of ${}^A N$. This vector space carries an irreducible representation, Π_ν^A , of

$D(G)$ defined by

$$\Pi_\nu^A(P_ab)|^Ag_j, {}^\nu v_i\rangle = \delta_{a,b^Ag_jb^{-1}}|b^Ag_jb^{-1}, D^\nu({}^Ax_l^{-1}b^Ax_j)^\nu v_i\rangle \quad (4.15)$$

where Ax_l is defined by ${}^Ag_l = b^Ag_jb^{-1}$. Notice that ${}^Ax_l^{-1}b^Ax_j$ is in AN . The gauge transformation b is “twisted” into the normalizer of the flux. It can be shown that these representations form a complete set of irreducible representations of $D(G)$. Any representation of $D(G)$ can be decomposed into a direct sum of these representations. In $|^Ag_j, {}^\nu v_i\rangle$, the conjugacy class labels the magnetic flux of the dyon, the representation of AN labels the electric charge. We can use the comultiplication to define the tensor product of representations of $D(G)$.

The state of an ordinary electrically charged particle is $|e, {}^\nu v\rangle$ where now, ν is an irreducible representation of G . The state of a single vortex in a group eigenstate is $|h, 1\rangle$ relative to some standard path, where the 1 is the trivial representation. It is found that $\Pi_1^{[h]} \otimes \Pi_1^{[h^{-1}]} = \Pi_\nu^{[e]} \oplus \dots$ where ν is a non-trivial representation of G and this is the Cheshire charge that a pair of vortex-anti-vortex can carry.^[16] If we consider the handle in a state $|a; b\rangle$ as a particle, it has magnetic flux $aba^{-1}b^{-1}$, and the operator h changes its state to $|hah^{-1}, hbh^{-1}\rangle$. The state of the whole handle has the same transformation properties under the quantum double as the state

$$|a, 1\rangle \otimes |b, 1\rangle \otimes |a^{-1}, 1\rangle \otimes |b^{-1}, 1\rangle . \quad (4.16)$$

For example, we can calculate the possible Cheshire charge of a handle by decomposing the tensor product $\Pi_1^{[a]} \otimes \Pi_1^{[b]} \otimes \Pi_1^{[a^{-1}]} \otimes \Pi_1^{[b^{-1}]}$. We must be careful about the meaning of the expression in (4.16). It is originally for the state of four vortices or dyons. In this case, it represents the state of a single handle. For example, we cannot apply the braiding operator to it.

5. Dyons on higher genus surfaces

For any surface, Σ , with n dyons on it, we can specify the state by choosing standard paths for the dyons and the handles and associating a vector in some representation of the quantum double for each path. One may expect that there is a correspondence between the multiplication of paths in the fundamental group and the tensor product of vectors in representations of the quantum double. However, the correspondence does not exist. To illustrate this, consider the product $C_1 C_1^{-1}$ in Fig. 2. It is trivial in the fundamental group. The state associated with it must be the vector in the trivial representation, but if the state associated with C_1 is $|h_1, 1\rangle$, the state associated with C_1^{-1} is $|h_1^{-1}, 1\rangle$. The tensor product $|h_1, 1\rangle \otimes |h_1^{-1}, 1\rangle$ transforms as a linear combination of charge eigenstates, not as the vector in the trivial representation. The reason why it does not work is that there is in general no “inverse” of a vector in any representation of the quantum double.*

This also occurs in ordinary spacetime. For example, consider QCD in $3 + 1$ dimensions. When we say that there are two units of red charge inside a closed surface, we mean that we have chosen the *outward normal* of the surface and after integrating the color electric field on the surface relative to this normal direction, we get two units of red charge. If we consider the product of the surface and itself with inward normal in the second homotopy group, and the tensor product of the corresponding charge, we run into the same difficulty as described above.

However, the tensor product does give us the combined state of two subsystems. In $3 + 1$ dimensions, we have to choose the outward normal (or inward normal) for both surfaces and determine the states corresponding to these surfaces. Then the state of the combined system is given by the tensor product. In our case of dyons, the orientations of the standard loops must be in the “same sense.” For example, if the states associated with C_1 and C_2 in Fig. 2 are $|h_1, 1\rangle$ and $|h_2, 1\rangle$ respectively, the state associated with $C_1 C_2$ is $|h_1, 1\rangle \otimes |h_2, 1\rangle$.

* I thank John Preskill for giving me this example.

For a compact surface, Σ_g , with n dyons, we choose the conventions in Fig. 8. The states of the dyons can be measured by charged particles and vortices traveling around C_i (see Appendix). We can denote the state where α_i maps to a_i , β_i maps to b_i and C_j maps to $|h_j, \nu_j v\rangle$ by

$$|a_1, b_1, \dots, a_g, b_g; h_1, \nu_1 v; \dots; h_n, \nu_n v\rangle . \quad (5.1)$$

A general state will be a linear combination of these.

The state must satisfy the relation discussed in section 3. This means that for the state

$$\sum_r k_r |a_i^{(r)}, b_i^{(r)}; h_j^{(r)}, \nu_j v_{(r)}\rangle , \quad (5.2)$$

where the k_r 's are constants, the tensor product

$$\begin{aligned} & \sum_r k_r |a_1^{(r)}, 1\rangle \otimes |b_1^{(r)}, 1\rangle \otimes |(a_1^{(r)})^{-1}, 1\rangle \otimes |(b_1^{(r)})^{-1}, 1\rangle \\ & \otimes \dots \otimes |a_g^{(r)}, 1\rangle \otimes |b_g^{(r)}, 1\rangle \otimes |(a_g^{(r)})^{-1}, 1\rangle \otimes |(b_g^{(r)})^{-1}, 1\rangle \\ & \otimes |h_1^{(r)}, \nu_1 v_{(r)}\rangle \otimes \dots \otimes |h_n^{(r)}, \nu_n v_{(r)}\rangle \end{aligned} \quad (5.3)$$

must transform as the trivial representation. (cf. (3.7))

We can also consider the motion of the dyons. Similar to the discussion in section 3, there is an action of the braid group on the states of the surface with dyons. For $B_n(\Sigma_g)$, the action of the σ 's are given by the $\mathcal{R}$ operator as discussed above.^[16] From (3.12), we also have

$$\begin{aligned} \alpha_j |a_j, b_j; h_1, \nu_1 v\rangle &= |a_j h_1 a_j h_1^{-1} a_j^{-1}, a_j h_1 a_j^{-1} h_1^{-1} b_j a_j h_1^{-1} a_j^{-1}\rangle \\ & \otimes \Pi_{\nu_1}^{[h_1]}(a_j) |h_1, \nu_1 v\rangle \\ \beta_j |a_j, b_j; h_1, \nu_1 v\rangle &= |a_j h_1, h_1^{-1} b_j h_1\rangle \otimes \Pi_{\nu_1}^{[h_1]}(h_1^{-1} b_j) |h_1, \nu_1 v\rangle . \end{aligned} \quad (5.4)$$

One can also check that this definition satisfies the relations (3.11). Notice that the action of the α 's and β 's cannot be written as the action of some element of the

quantum double because the flux of α and β are not just conjugated. This represents the fact that if we do see the internal structure of the handle, its state is not in the Hilbert space of the states of a particle. If every dyon does not carry electric charge, the representations ν_i are trivial. All formulae here then reduce to the corresponding formulae in section 3.

Let us consider an example. Suppose the group G is the quaternion group $Q = \{\pm 1, \pm i, \pm j, \pm k\}$. There are five conjugacy classes: $\{1\}$, $\{-1\}$, $\{\pm i\}$, $\{\pm j\}$ and $\{\pm k\}$. And there are four one-dimensional irreducible representations: the trivial representation 1 and $1_x, 1_y, 1_z$ where in 1_x , say, $\pm 1, \pm i$ are represented by 1 and the others are represented by -1 . There is also a two-dimensional irreducible representation. Notice that the normalizer of -1 is the whole group. So, if a dyon has flux -1 , its electric charge can be labeled by representation of the whole group.

Assume the space is a torus and there are two vortices and one charged particle. A possible state is

$$|v\rangle = \frac{1}{2}(|i, j; k, 1; k, 1; 1, 1_x\rangle + |i, -j; -k, 1; -k, 1; 1, 1_x\rangle - | -i, j; -k, 1; -k, 1; 1, 1_x\rangle - | -i, -j; k, 1; k, 1; 1, 1_x\rangle) . \quad (5.5)$$

In each term, the first two factors label the flux carried by α and β ($\pm i$ and $\pm j$). The third and fourth factors are the flux ($\pm k$) and the charge (trivial) of the first vortex. The next two factors have the same meaning. The final two are the trivial flux (1) and the charge (1_x) of the charged particle. The state of the handle and the states of the vortices are entangled but if we consider them as a whole, they are in the state $|1, 1_x\rangle$, so together with the charged particle, they satisfy (5.3).

If the first vortex winds around β , by (5.4), we have

$$\begin{aligned} \beta|v\rangle &= \frac{1}{2}(| -j, -j; k, 1; k, 1; 1, 1_x\rangle + |j, j; -k, 1; -k, 1; 1, 1_x\rangle \\ &\quad - | -j, -j; -k, 1; -k, 1; 1, 1_x\rangle - |j, j; k, 1; k, 1; 1, 1_x\rangle) \\ &= \frac{1}{2}(| -j, -j\rangle - |j, j\rangle) \otimes (|k, 1\rangle \otimes |k, 1\rangle - | -k, 1\rangle \otimes | -k, 1\rangle) \otimes |1, 1_x\rangle . \end{aligned} \quad (5.6)$$

Now, the first factor in the above tensor product is the state of the handle, the

second factor is the state of the two vortices and the last factor is the state of the charged particle. The handle carries magnetic flux -1 and Cheshire charge 1_y . The two vortices together carry flux -1 and charge 1_z . There is magnetic flux transfer between the handle and the pair of vortices.

What will happen if a charged particle winds around a loop of a handle? As we have seen in the beginning of section 4, if the handle is in flux eigenstate, the state of the particle will be transformed by the flux of the loop and the state of the handle will remain the same. If the handle is in some linear combination of flux eigenstates, something interesting will happen. For example, let the state of the charged particle be $|v\rangle$, and assume initially the state of the handle is $\frac{1}{\sqrt{|G|}} \sum_{b \in G} |e, b\rangle$. Then the charge bounded by α is trivial. If the charged particle winds around β , then

$$|v\rangle \otimes \frac{1}{\sqrt{|G|}} \sum_{b \in G} |e, b\rangle \rightarrow \frac{1}{\sqrt{|G|}} \sum_{b \in G} |D(b)v\rangle \otimes |e, b\rangle . \quad (5.7)$$

If we now introduce a h vortex, and wind it around α , the state changes to $\frac{1}{\sqrt{|G|}} \sum_{b \in G} |D(b)v\rangle \otimes |e, bh^{-1}\rangle$, then the interference term is proportional to

$$\begin{aligned} & \sum_{b, b' \in G} \langle e, b' | \langle D(b')v | D(b)v \rangle | e, bh^{-1} \rangle \\ &= \sum_{b, b' \in G} \langle v | D(b'^{-1}b) | v \rangle \delta_{b', bh^{-1}} \\ &= \sum_{b \in G} \langle v | D(h) | v \rangle . \end{aligned} \quad (5.8)$$

The charge bounded by α is v and the flux of α is identity. We see that the charge of the particle is transferred to the ingoing mouth of the handle. However, the state of the particle entangles with the state of the handle and can no longer be specified by a single vector. This kind of charge transfer between mouths of wormholes or handles and charged particles also occurs in $3 + 1$ dimensions and for continuous gauge groups.^[21]

6. Conclusion

We have argued that in $2 + 1$ dimensions, non-trivial topology, the handle, can carry magnetic flux classically. If the unbroken gauge group is finite, we can actually assign arbitrary group elements to the two non-equivalent loops associated to the handle. Semi-classically, the state of a handle can be specified by the flux of the two non-trivial loops. It can also carry Cheshire charge. On the other hand, a general particle will fall into representations of the quantum double, an algebra constructed from the gauge group. We have also explained the physical meanings of the elements of the quantum double, the comultiplication and the braiding operator $\mathcal{R}$. If the surface is compact, there is a relation between the generators of the fundamental group of the space. This relation restricts the possible flux and charges of the handles and the dyons on that surface. If the surface is non-compact, no such relation exists.

There is topological interaction between the dyons and the handles. The motion of the dyons is described by the braid group of the surface. Then, the topological interaction can be described by an action of the braid group on the states of the handles and the dyons. This action, and hence the topological interaction, can be completely determined by the path tracing method explained in section 2. A similar classical analysis in $3 + 1$ dimensions for cosmic strings has been done by Brekke et. al.,^[22] and the classification of dyons has been generalized to theories with Chern-Simons terms.^[23]

APPENDIX

In this appendix, we will recall how to measure the flux of a beam of identical vortices using charged particles^[8] and how to measure the charge of a beam of identical charged particles using vortices.

Assume that we have a beam of identical vortices with unknown flux, h , and we have charged particles in any desired states. We can send the charged particles in a particular state around the vortices and then observe the interference patterns. If the

state of the particles is $|v\rangle$ in some representation ν , the interference gives us

$$\langle v|D^{(\nu)}(h)|v\rangle . \quad (\text{A.1})$$

If we replace $|v\rangle$ by $|v\rangle + \lambda|w\rangle$ in the above equation where λ is an arbitrary complex number and subtract $\langle v|D^{(\nu)}(h)|v\rangle + \langle w|D^{(\nu)}(h)|w\rangle$ from it, we have

$$\lambda\langle v|D^{(\nu)}(h)|w\rangle + \lambda^*\langle w|D^{(\nu)}(h)|v\rangle . \quad (\text{A.2})$$

Put λ to be i and 1 in succession, we know the values of two expressions. A linear combination of them gives us

$$\langle v|D^{(\nu)}(h)|w\rangle . \quad (\text{A.3})$$

We can now determine $\langle v|D^{(\nu)}(h)|w\rangle$ for arbitrary $|v\rangle, |w\rangle$ and ν and hence the matrix representation of h . If we choose ν to be some faithful representation, we can determine h .

Now assume that we have a beam of charged particles in some unknown state, $|v\rangle$, in some unknown irreducible representation, ν , and we have vortices with any desired flux. We also assume that $\langle v|v\rangle = 1$. Then a similar interference experiment will give us

$$\langle v|D^{(\nu)}(h)|v\rangle \quad (\text{A.4})$$

for arbitrary h . Because ν is irreducible, the vectors, $D^{(\nu)}(h)|v\rangle$ for $h \in G$, will span the whole representation space. We know the inner products of these vectors because $\langle D^{(\nu)}(h_1)v|D^{(\nu)}(h_2)v\rangle = \langle v|D^{(\nu)}(h_1^{-1}h_2)|v\rangle$. By Gram-Schmidt's orthogonalization, we can form a basis, $\{|e_i\rangle : i = 1, \dots, d\}$, such that $|e_1\rangle = |v\rangle$ and

$$|e_i\rangle = \sum_{h \in G} c_h^i |D^{(\nu)}(h)v\rangle . \quad (\text{A.5})$$

Notice that the coefficients, c_h^i , depend only on the numbers $\langle v|D^{(\nu)}(h')|v\rangle$. We also have $|D^{(\nu)}(h)v\rangle = \sum_i b_i^h |e_i\rangle$ for some coefficients b_i^h .

Now we have a basis, so we can calculate the character of the representation and hence determine the representation itself.

Suppose that there is another vector, $|w\rangle$, in the same representation space such that for all h in G , $\langle w|D^{(\nu)}(h)|w\rangle = \langle v|D^{(\nu)}(h)|v\rangle$. We are going to prove that $|w\rangle$ is equal to $|v\rangle$ up to a phase. If this can be done, we can uniquely determine the state of the beam of charged particles by only sending vortices around them and observing the interference pattern.

Let $|e'_i\rangle = \sum_{h \in G} c_h^i |D^{(\nu)}(h)w\rangle$. Since the coefficients, c_h^i , depend only on the numbers $\langle v|D^{(\nu)}(h')|v\rangle = \langle w|D^{(\nu)}(h')|w\rangle$, $|e'_i\rangle$ form a basis. Then there is an operator L such that $|e'_i\rangle = L|e_i\rangle$. We claim that $LD^{(\nu)}(h) = D^{(\nu)}(h)L$ for all h in G . First of all, we have $|w\rangle = |e'_1\rangle = L|e_1\rangle = |v\rangle$. Then, $D^{(\nu)}(h)L|v\rangle = D^{(\nu)}(h)|w\rangle = \sum_i b_i^h |e'_i\rangle = \sum_i b_i^h L|e_i\rangle = L \sum_i b_i^h |e_i\rangle = LD^{(\nu)}(h)|v\rangle$. We have

$$\begin{aligned} D^{(\nu)}(h)L|e_i\rangle &= D^{(\nu)}(h)|e'_i\rangle = \sum_{h' \in G} c_{h'}^i D^{(\nu)}(hh')|w\rangle \\ &= \sum_{h' \in G} c_{h^{-1}h'}^i D^{(\nu)}(h')|w\rangle = \sum_{h' \in G} c_{h^{-1}h'}^i D^{(\nu)}(h')L|v\rangle \\ &= L \sum_{h' \in G} c_{h^{-1}h'}^i D^{(\nu)}(h')|v\rangle = LD^{(\nu)}(h)|e_i\rangle. \end{aligned} \tag{A.6}$$

This proves the claim. Since ν is irreducible, by Schur's lemma, L is the product of a constant and the identity operator and $|w\rangle = e^{i\theta}|v\rangle$ because $\langle w|w\rangle = \langle v|v\rangle = 1$.

For a beam of dyons, we can first measure the magnetic flux by sending charged particles around them. After we know the flux, we can measure their charge by using vortices with flux which commutes with the flux of the dyons. Then, we completely determine the state of the dyons in some representation of the quantum double.

This analysis can be generalized to the measurement of a single particle or particles in a reducible representation (with some limitation).^[8]

Acknowledgments

I would like to thank Hoi-Kwong Lo and especially John Preskill for very useful discussions.

REFERENCES

1. J. Preskill, Vortices and monopoles, *in* Architecture of the fundamental interactions at short distances, ed. P. Ramond and R. Stora (North-Holland, Amsterdam, 1987).
2. Y. Aharonov and D. Bohm, Phys. Rev. 119 (1959) 485.
3. A. S. Schwarz, Nucl. Phys. B208 (1982) 141.
4. L. Krauss and F. Wilczek, Phys. Rev. Lett. 62 (1989) 1221.
5. M. Alford, K. Benson, S. Coleman, J. March-Russell and F. Wilczek, Phys. Rev. Lett. 64 (1990) 1632; Nucl. Phys. B349 (1991) 414.
6. J. Preskill and L. Krauss, Nucl. Phys. B341 (1990) 50.
7. M. Bucher, Nucl. Phys. B350 (1991) 163.
8. M. Alford, S. Coleman and J. March-Russell, Nucl. Phys. B351 (1991) 735.
9. M. Alford and J. March-Russell, Nucl. Phys. B269 (1992) 276.
10. M.G. Alford, K.-M. Lee, J. March-Russell and J. Preskill, Nucl. Phys. B384 (1992) 251.
11. M. Bucher, H.-K. Lo and J. Preskill, Nucl. Phys. B386 (1992) 3.
12. M. Bucher, K.-M. Lee and J. Preskill, Nucl. Phys. B386 (1992) 27.
13. Y. Ladegaillerie, Bull. Sci. Math. 100 (1976) 255.
14. T.D. Imbo and J. March-Russell, Phys. Lett. B252 (1990) 84.
15. R. Dijkgraaf, V. Pasquier and P. Roche, Nucl. Phys. B (Proc. Suppl.) 18B (1990) 60.
16. F.A. Bais, P. van Driel and M. de Wild Propitius, Phys. Lett. B280 (1992) 63.

17. See, e.g., H.M. Farkas and I. Kra, *Riemann Surfaces*, Springer-Verlag 1980;
O. Forster, *Lectures on Riemann Surfaces*, Springer-Verlag 1981.
18. P. Nelson and A. Manohar, *Phys. Rev. Lett.* 50 (1983) 943; A. P. Balachandran,
G. Marmo, N. Mukunda, J. S. Nilsson, E. C. G. Sudarshan and F. Zaccaria,
Phys. Rev. Lett. 50 (1983) 1553; P. Nelson and S. Coleman, *Nucl. Phys.* B237
(1984) 1; A. P. Balachandran, F. Lizzi and V. G. Rodgers, *Phys. Rev. Lett.* 52
(1984) 1818; P. A. Horváthy and J. H. Rawnsley, *J. Math. Phys.* 27 (1986) 982.
19. P. Bantay, *Phys. Lett.* B245 (1990) 477; *Lett. Math. Phys.* 22 (1991) 187.
20. For a review of quantum doubles and quantum groups, see e.g., T. Tjin, *Int. J.*
Mod. Phys. A7 (1992) 6175.
21. H.-K. Lo and K.-M. Lee, *Wormholes and Charge Conservation*, Caltech preprint
CALT-68-1880 (1993).
22. L. Brekke, H. Dykstra, S.J. Hughes and T.D. Imbo, *Phys. Lett.* B288 (1992)
273.
23. F. A. Bais, P. van Driel and M. de Wild Propitius, *Nucl. Phys.* B393 (1993)
547.